



Original Research Article

Correlation between iron deficiency anemia and HbA1C levels in type 2 diabetes mellitus

K Sumathi¹, Gayathri Dilliraj^{1,*}, B Shanthi¹, V S Kalai Selvi¹, A Jamuna Rani¹¹Dept. of Biochemistry, Sree Balaji Medical College and Hospital, (Bharath University), Chennai, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 28-07-2020

Accepted 18-08-2020

Available online 28-09-2020

Keywords:

Iron deficiency anaemia status
Glycosylated Haemoglobin level (HbA1c)
Diabetes mellitus

ABSTRACT

Introduction: The HbA1c levels, in general reflects the diabetic patients Glycaemic control status over the previous 3 months. HbA_{1c} concentrations may be increased by Anaemia due to iron deficiency, independent of glycaemia in diabetic patients. Henceforth this study serves to find if there is a positive correlation between the iron deficiency anaemia status & the HbA_{1c} levels amidst the diabetic population.

Materials and Methods: Cases-100 iron-deficiency anaemia patients with diabetes having controlled plasma glucose levels, Controls -100 non-anaemic diabetic individuals. The study was done in the premises of Sree Balaji Medical College & Hospital. HbA_{1c} levels was measured by ion exchange chromatography, ferritin by particle enhanced turbidometric immunoassay method, fasting plasma glucose by GOD - POD method.

Results: We found that HbA_{1c} was elevated (7.3±0.9) in iron-deficient diabetic individuals compared to normal level (5.4±0.6) in controls.

Conclusion: The study showed that there was a positive correlation between the iron deficiency anaemia status and increased HbA_{1c} levels in the controlled diabetic population. Hence forth, the clinical significance of evaluating the iron deficiency anaemia status in diabetic patients has been proved to be significant in planning their further course of prompt diabetic management.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Among the various known anaemia's in India, Iron deficiency anaemia is more commonly encountered in both the urban and the rural forms of our society. The most common cause of this is under nutrition particularly amidst females. 50% of anaemia is attributed to iron deficiency, worldwide. The iron status in our body is precisely and accurately predicted by the ferritin levels (iron storage form).¹ Various previous studies have proved that, depleted iron status, tend to have a link with the deranged (increased) glycation of haemoglobin (HbA_{1c}) levels. HbA_{1c} is suggestive of the glycated haemoglobin content that is widely used as an indicator of a patient's glycaemic status over the previous 3 months.² HbA_{1c} levels act as the most accurate indicator of the individual's

diabetic status globally. According to the standards set by the American Diabetes Association Guidelines, HbA_{1c} levels below 7% in an diabetic patient, is termed as good control, that keeps them safe from the developing various life threatening micro vascular complications associated with diabetes.³ Certain haematological factors like the haemolytic anaemia's,⁴ hemoglobinopathies, and acute and chronic blood loss also have a significant impact on the HbA_{1c} levels besides pregnancy and uraemia. HbA_{1c} levels in the body may also be influenced/alterd by various other minor factors like Vitamin B12 & folate levels in addition to the iron deficiency anaemia status. Henceforth in this study we aimed to correlate the significance of estimating the iron deficiency status along with the HbA_{1c} levels, in our study's diabetic population.

* Corresponding author.

E-mail address: dr.gayathridilliraj@gmail.com (G. Dilliraj).

2. Materials and Methods

The study was done in the Medicine department in association with the Central laboratory, in the premises of, Sree Balaji Medical College & Hospital, Chromepet Chennai. The study chiefly consisted of 200 subjects consisting of 100 cases and 100 controls:

1. Cases-100 iron deficiency anaemia patients with diabetes having controlled plasma glucose levels,
2. Controls -100 non anaemic diabetic individuals.
3. Ion exchange chromatography – used to measure the HbA_{1c} levels.
4. Particle enhanced turbidometric immunoassay method – used to measure ferritin levels
5. GOD – POD method – to measure fasting blood glucose levels.
6. Haemoglobin, peripheral smear examination, red cell indices, and medical history were all recorded.

3. Results

Table 1:

Parameters	Patients	Controls
Haemoglobin	6.0±3.1	13.0±0.5
Serum ferritin(ng/ml)	6.5±3.5	235±75.5
HbA _{1c} %	7.3±0.9	5.4±0.6

7.3% was the mean HbA_{1c} levels in the anaemic patients, while that in the controls was 5.4% which is significant. ($p < 0.01$).

4. Discussion

This is a typical case control study organised and conducted in the premises of Sree Balaji Medical College & Hospital. The most common fraction of HbA₁ is HbA_{1c}. HbA_{1c} is predominantly formed by the interaction of Glucose with the N-terminal valine groups of both the beta chains, to form an aldimine linkage that ultimately realigns itself to form the stable ketoamine link of HbA_{1c}.^{5,6} HbA_{1c} is a globally acknowledged significant criteria, for diagnosing, treating and to overview the prognosis of the diabetic patients, in accordance with the American Diabetic Association. HbA_{1c} levels in the body may also be influenced/alterd by various other common haematological conditions like;⁷⁻⁹ haemolytic anaemia's, acute and chronic blood loss, etc. Pregnancy and uraemia also have a significant impact on HbA_{1c} levels.^{2,10,11}

For decades, both the urban and the rural population of India have encountered varying degrees of Iron deficiency anaemia. In accordance with various global studies it was proved that iron therapy proved to be beneficial in maintaining a well-controlled glycated haemoglobin level.

There was a significant decrease in HbA_{1c} levels after iron therapy even in the non-diabetic population to some extent.^{2,12,13} In the erythrocytes as the age advances, the HbA₁ levels also tend to increase.¹⁴ This age related haemoglobin glycation process is of an irreversible type. As we age there is also a depletion in the production of RBC that is concomitantly accompanied by an increased age of circulating red cells with increased HbA₁ levels.¹⁴

In our study it was scientifically proved that the elevated HbA_{1c} levels were caused significantly due to the iron deficiency status. Although the HbA_{1c} levels are generally higher in the diabetic patients, patients with controlled plasma glucose levels for 3 months are generally expected to have HbA_{1c} level below 6.5%.⁴

The results showed that, the HbA_{1c} levels were significantly elevated in iron-deficient anaemic individuals with FPG less than 130. The importance of HbA_{1c} as an important prognostic marker has been established by various other studies carried out in diabetic individuals.^{10,15,16} From this study the positive correlation between iron deficiency anaemia and elevated HbA_{1c} levels have been scientifically proved, that should always be considered before altering diabetic treatment regimen. Iron is stored chiefly in the body in the form of ferritin, its storage form that is highly suggestive, of the iron load status in the body.¹

Increased RBC life span with concomitantly increased HbA_{1c} levels and decreased ferritin levels are encountered in most of the cases of iron deficiency anaemia. Previous studies have enhanced the fact, that with iron therapy there tends to be a clinically significant decrease in the elevated HbA_{1c} values of diabetic patients with iron deficiency anaemia.² Iron therapy also showed improvement (i.e. decrease) in the previously increased HbA_{1c} status of pregnant women with diabetes and diabetic CKD (chronic kidney disease) patients who were suffering with iron deficiency anaemia.^{10,11} Thus it was concluded that the HbA_{1c} levels in diabetic patients was generally elevated when they had iron deficiency status, when compared with iron-sufficient controls. HbA_{1c} continues to be the most simple yet a significant marker for glycaemic control estimation. Thus the various other factors that tend to elevate the HbA_{1c} levels, should also be excluded and accounted for, before planning the further course of management in a diabetic patient.

5. Conclusion

Henceforth this study, scientifically and statistically proved the positive correlation between the iron deficiency anaemia status and the increased HbA_{1c} levels in the controlled diabetic individuals. Thereby insisting the diabetic population to periodically check their iron status too, that has the potency to influence their HbA_{1c} parameter used to assess the degree of their glycaemic control before

planning the course of management for the diabetes cases.

6. Source of Funding

None.

7. Conflict of Interest

None.

8. Acknowledgement

All the participants/subjects were explained in detail about the purpose of the study. And consent forms were obtained from all of them.

References

- Wyck DBV, Stivelman JC, Ruiz J, Kirilin LF, Katz MA, Ogden DA. Iron status in patients receiving erythropoietin for dialysis-associated anemia. *Kidney Int.* 1989;35(2):712–6.
- Sherwani SI, Khan HA, Ekhzaimy A, Masood A, Sakharkar MK. Significance of HbA1c Test in Diagnosis and Prognosis of Diabetic Patients. *Biomark Insights.* 2016;11:95–104.
- Qaseem A, Wilt TJ, Kansagara D, Horwitch C, Barry MJ, Forciea MA. Haemoglobin A1C Targets for Glycaemic Control with Pharmacologic Therapy for Non pregnant Adults with Type 2 Diabetes Mellitus: A Guidance Statement Update from the American College Of Physicians. *Ann Intern Med.* 2018;168(8):569–76.
- Standards of Medical Care in Diabetes-2018 Abridged for Primary Care Providers. *Clin Diabetes.* 2018;36(1):14–37.
- Bunn HF, Haney, Kenneth P. Further Identification of the Nature and Linkage of the Carbohydrate in Hba1c. *Biochem Biophys Res Commun.* 1975;67(1):103–9.
- Bonora E, Tuomilehto J. The Pros and Cons of Diagnosing Diabetes With A1C. *Diabetes Care.* 2011;34(Supplement_2):S184–90.
- Sultanpur CM, Deepa K, Kumar SV. Comprehensive Review on Hba1c in Diagnosis of Diabetes Mellitus. *Intl J Pharm Sci Rev Res.* 2010;3(2):119–22.
- Kumar PR, Bhansali A, Ravikiran M, Bhansali S, Dutta P, Thakur JS. Utility of Glycated Hemoglobin in Diagnosing Type 2 Diabetes Mellitus: A Community-Based Study. *J Clin Endocrinol Metab.* 2010;95(6):2832–5.
- Sarmah D, Sharma B. Importance and Status of HBA1C in T2DM and its Indian Perspective. *Asian J Biomed Pharm Sci.* 2012;2(12):1–10.
- Sinha N. HbA1c and factors other than diabetes mellitus affecting it. *Singapore Med J.* 2010;51(8):616–22.
- Rafat D, Rabbani TK, Ahmad J, Ansari MA. Influence of iron metabolism indices on HbA1c in non-diabetic pregnant women with and without iron-deficiency anemia: Effect of Iron supplementation. *Diabetes Metab Syndr Clin Res Rev.* 2012;6(2):102–5.
- Kumar M, Nutakki G. A prospective study of effect of iron deficiency anaemia on HbA1c levels in nondiabetics. *IAIM.* 2017;4(12):137–46.
- Lodh M, Mukhopadhyay R, Kumar B. A Case of Inappropriately High Glycated Hemoglobin. *Indian J Clin Biochem.* 2015;30(2):234–7.
- Zhang C, Rawal S. Dietary iron intake, iron status, and gestational diabetes. *Am J Clin Nutr.* 2017;106(Supplement 6):1672S–80.
- Christy AL, Manjrekar PA, Babu P, Hegde A, Rukmini MS. Influence of Iron Deficiency Anemia on Haemoglobin A1C Levels in Diabetic Individuals with Controlled Plasma Glucose Levels. *Iran Biomed J.* 2014;18(2):88–93.
- Luo J F Li JH, Zhang HD. Red blood cell lifespan shortening in patients with early stage chronic kidney disease. *Kidney Blood Press Res.* 2019;44:1158–1165.

Author biography

K Sumathi Associate Professor

Gayathri Dilliraj Tutor

B Shanthi Professor and HOD

V S Kalai Selvi Professor

A Jamuna Rani Associate Professor

Cite this article: Sumathi K, Dilliraj G, Shanthi B, Selvi VSK, Rani AJ. Correlation between iron deficiency anemia and HbA1C levels in type 2 diabetes mellitus. *Int J Clin Biochem Res* 2020;7(3):400–402.