

## Altered erythrocyte surface electric charge in $\beta$ -thalassemia: By Alcian Blue Dye binding

Dhiraj J. Trivedi<sup>1,\*</sup>, Vishwaraj PK<sup>2</sup>

<sup>1</sup>Professor, <sup>2</sup>UG Student, Dept. of Biochemistry, SDM College of Medical Sciences & Hospital, Dharawad, Karnataka

**\*Corresponding Author:**

Email: dhiraj99trivedi@gmail.com, trivedi9962dhiraj@yahoo.co.uk

### Abstract

**Aim:** To evaluate erythrocyte surface electric charge in  $\beta$ -thalassemia by using Alcian Blue dye binding.

**Objective:** Estimate the binding of AB dye to Sialic acid on the surface of thalassemia and normal, healthy erythrocytes. Compare negative charge due to Sialic acid on the surface of erythrocyte of thalassemia with healthy control subjects.

**Result:** The study was performed on 60 thalassemia subjects (39 male, 21 female, mean age 8.3years) and 30 age and sex matched normal control subjects (17 male, 13 female, mean age 6.8years). Hematological parameters analyzed on beta thalassemia subjects' shows significant difference among control and Thalassemia group. Levels of HbA<sub>2</sub> showed a marginal difference (p=0.041). Cationic dye, Alcian Blue, in PBS (pH7.4) was used as a reagent to demonstrate binding of the dye to Sialic acid. Results of dye bound to thalassemic erythrocytes was significantly less (p<0.05) as compared to binding on erythrocytes of the control group.

**Conclusion:** We believe, our data support the perception that there is decreased negative charge on the erythrocyte membrane in thalassemia reflected by cationic Alcian Blue dye binding.

**Keywords:** Thalassemia, Erythrocyte, Alcian Blue Dye, RBC Membrane, Sialic acid.

### Introduction

The clinical significance of Glycobiology has become the focus of today's research, as its role in the development, regulation and progression of the disease is slowly but, surely recognized. Sialic acid, a component of glycoprotein, plays a central role in the biochemical function of humans. Elevated levels of Sialic acid in metabolic diseases, infectious diseases, inflammatory diseases and carcinogenesis are well established. Altered sialoglycoprotein and Sialic acid on erythrocyte membrane has also been reported in various diseases like, Thyroid disorder, Diabetes, Jaundice, Nephrotic disease, glomerular disease, Alcoholism, cardiac disease, RA and various forms of malignant transformation. In addition, some physiological process may alter Sialic acid levels in plasma.<sup>(1)</sup>

Thalassemia, a genetic disease associated with abnormal hemoglobin, a defect in the beta globin chain cluster. The Beta thalassemia carrier rate in India is around 3-7% with higher frequency in certain ethnic groups.<sup>(2)</sup> Thalassemia is a form of transfusion dependent hemolytic type of anemia accompanied by iron overload and hemochromatosis as a side effect of repeated blood transfusion. Microcytic, hypochromic, anisocytic red cells with basophilic stippling and target cell appearance with increased osmotic fragility is a hematological hallmark of thalassemic erythrocytes.<sup>(3)</sup> Presence of unstable hemoglobin,<sup>(4)</sup> Iron overload,<sup>(5,6)</sup> Increased Reactive Oxygen Species along with reduced red cell Glutathione<sup>(7,8)</sup> and denatured membrane protein, loss of Sialic acid from glycol-conjugates of membrane<sup>(9)</sup> are some biochemical defects reported in thalassemia.

Morphological alterations and maldistribution of glycoprotein reported in thalassemic red cells are associated with damage of the red cell membrane skeleton.<sup>(10)</sup>

Recently, interest of Glycobiology has been shifted towards characterizing membrane flaw in hemoglobinopathies. Though such defect is not a primary cause, but it has some impact on survival of erythrocytes in these genetic diseases. The structure of Membrane micro domain or raft's contribution towards physical properties of live cell membranes continues to be clarified. Sialic acid a component of glycoprotein contributes almost all the negative charge present on the surface of erythrocyte membranes. Such high negative charge believed to suppress erythrocyte aggregation, but when removed results in altered rheological properties and causes aggregation of erythrocyte. It marks aging of circulating erythrocyte and causes early clearance from circulation. Thus Sialic acid has becomes a useful parameter to study the impact of genetic disease on modifications of cell surfaces micro domain.

Cationic dye Alcian Blue (AB) is a routine histochemical stain used for staining sialylated glycoalkal of cell. AB dye specifically binds with sialylated glycoalkal of cells.<sup>(11)</sup> It has been used to compare Sialic acid and in turn map electric charge on erythrocyte in various diseases.<sup>(12-14)</sup> Gambaro G et al<sup>(15)</sup> and Baba Y et al<sup>(16)</sup> have reported altered erythrocyte charge in diabetes mellitus. Levin et al<sup>(17)</sup> reported reduced binding of cationic AB dye to erythrocyte in children with minimal change nephrotic syndrome. D Yavuz et al<sup>(18)</sup> found reduced binding of AB in hypertensive patients. Changes have also been

recognized in PAS pattern of beta thalassemia RBC membrane protein electrophoresis. In view of existing research, we undertook a study to evaluate the surface charge on thalassemia erythrocyte membrane by using Alcian Blue dye binding.

### Material and Methods

The present study was performed at the Department of Biochemistry, of SDM College of Medical Sciences, Dharwad, Karnataka, as a part of ICMR-STS-2016 project. The Proposal was approved by the institutional ethics committee with a caution that the identity of the patient and results should be kept confidential.

The study was performed on 90 subjects of both genders between 5 and 21 years age group, living in and around Dharwad town in North Karnataka. The study population was categorized into two groups. Group A- consisted of sixty Thalassemia patients, diagnosed based on red cell indices (Hb < 10.0 gm/dl, RBC count >  $5 \times 10^6$  /cmm, RDWI < 17%, MCV < 80fl, peripheral smear showing Microcytic, Hypochromic RBC with target cells and/or basophilic stippling as suggested by Shine and Lal<sup>(19)</sup> and showing HbA<sub>2</sub> > 3.5% by Hemoglobin electrophoresis.<sup>(20)</sup> Group B- comprised of thirty age and sex matched children visiting pediatric OPD of our hospital who are apparently healthy and without any history of blood abnormalities. Objective of our study was to estimate Sialic acid on the surface of erythrocyte in thalassemia and normal, healthy subjects by Alcian blue dye binding method and compare negative charge due to Sialic acid on the surface of erythrocyte of both above groups.

Written informed consent was obtained from each participant or family member included in the research study

#### Preparation of Alcian Blue (cationic dye) solution:

As suggested by Winkel JGJ et al,<sup>(14)</sup> 5mg of Alcian Blue dye was completely dissolved in 100microliter of absolute ethanol and diluted to 10ml with Phosphate Buffered saline (PBS), pH7.4 containing 25mmol MgCl<sub>2</sub>. The final concentration of the AB dye was 500microgram/ml. Every time a fresh dye solution was prepared and extinction of dye solution was measured on the spectrophotometer (UV1700 Simadzu) at 650nm. Only the dye solution with extinction reading between 0.800 and 1.130 was accepted for further test.

**Collection of Blood and preparation of washed red blood cells:** After taking all aseptic precautions 2ml blood was collected from an anti-cubital vein in a tube

containing 250microlit of 3.8% trisodium citrate solution. The blood sample was centrifuged at 3000 rpm for 10 min to obtain the packed erythrocyte fraction. Plasma was discarded and packed erythrocytes were washed thrice with Phosphate Buffered saline (PBS).

**Cell count and Red Blood cell indices:** Fifty microliter of packed and washed erythrocyte were transferred to 10.0 ml of PBS. It was divided into two portions of 5ml each. A first aliquot of 5ml suspension was subjected for blood indices and RBC count on Sysmax XN 1000 – Transasia blood counter in hematology laboratory of our hospital.

**Cationic dye Alcian Blue binding to erythrocyte:** To the second aliquot of 5ml erythrocyte suspension, 500microliter of cationic dye, Alcian Blue (AB) was added. The suspension was incubated for 30min at 37 °C. After incubation for 30 min, erythrocytes were removed by centrifugation and leftover concentration of AB dye in solution was estimated by measuring optical density reading at 650nm on UV1700 Shimadzu Spectrophotometer. The amount of AB dye bound to erythrocyte was calculated. The final result was expressed as amount of AB dye bound per 10<sup>6</sup> erythrocyte.

**Hemoglobin electrophoresis:** 250microliter of washed packed erythrocyte were used for preparation of hemolysate which was subjected to alkaline hemoglobin electrophoresis on agarose gel supplied by Helena Co (Alere) to estimate concentrations of Hemoglobin fractions. Electrophoretogram was quantitated on densitometer using Platinum Ver 3.0 Software supported by Helena Co.

The final result were expressed in terms of dye bound per 10<sup>6</sup> red blood cells. Results were compared in patients and control groups and presented as mean + SD. Data was analyzed for Students 't' test within 95% CI by SPSS statistical software. P value <0.05 was considered as significant.

### Observations and Results

The study was performed on 60 thalassemia subjects (39 male, 21 female, mean age 8.3years) and 30 age and sex matched normal control subjects (17 male, 13 female, mean age 6.8years).

To support the selection of B thalassemia subjects hematological parameters were analyzed for whole blood collected from control and Thalassemia group. Hemoglobin A<sub>2</sub> levels were used as confirmatory parameter in diagnosis.

**Table 1: Hematological parameter in Control and Thalassemia Group**

	Control Group	Thalassemia Group	P value
Hemoglobin gm%	13.9 0.9 (11.8 – 15.7)	10.39 0.69 (5.3 – 17.6)	< 0.01
RBC count 10 <sup>6</sup> /cmm	4.58 0.36 (3.88 – 5.92)	5.56 0.4 (2.04 – 7.89)	< 0.001
MCV fl	86.2 4.7 (79.5 – 99.6)	60.11 3.49 (52.7 – 85.3)	< 0.01
MCH Pg	30.1 1.6 (25.4 – 35.8)	20.76 4.69 (16.5 27.2)	< 0.05
RCWD %	14.2 0.98 (9.9 – 34.4)	10.76 1.38 (9.3 – 31.6)	< 0.05
Hemoglobin A2 %	2.3 0.6 (1.8 – 3.3)	5.05 1.30 (3.6 – 8.5)	= 0.041

Result of control and Thalassemia groups for Hemoglobin, RBC count, MCV, MCH, RCWD and HbA2 from Table 1, when worked out for student t test showed a significant difference in hemoglobin ( $p < 0.01$ ), RBC count ( $p < 0.001$ ) MCV ( $p < 0.01$ ), MCH ( $P < 0.05$ ) and RCWD ( $p < 0.05$ ) between two groups. HbA2 showed a marginal difference ( $p = 0.041$ ) between both the groups.


Cationic dye, Alcian Blue, in PBS (pH7.4) was used as a reagent to demonstrate binding of the dye to Sialic acid moieties on the surface of erythrocytes from thalassemia and control groups.

**Table 2: Cationic Alcian Blue dye binding in Control and Thalassemia Group**


	Control Group Male n= 17(56.6%)	Control Group Female n= 13(43.3%)	Thalassemia Group Male n= 39(65%)	Thalassemia Group Female n= 21(35%)	P value
ng AB dye Unbound in sol /10 <sup>6</sup> RBC	405.2 ± 9.8	398.1 ± 6.2	301.2 ± 5.2	313.6 ± 8.3	
ng AB Dye Bound/10 <sup>6</sup> RBC	116.4 ± 1.8	129.5 ± 2.2	95.2 ± 0.9	98.7 ± 1.3	
ng AB dye bound to 10 <sup>6</sup> RBC (Average)	127.9		95.1		<0.05

Binding of cationic dye Alcian Blue on erythrocyte in control and thalassemia group is shown in Table 2. In our study, Mean value of dye bound to male erythrocytes of control group was  $116.4 \pm 1.8$  ng/10<sup>6</sup> RBC and female erythrocytes of control group was  $129.5 \pm 2.2$  ng/10<sup>6</sup> RBC. Whereas the mean value of dye bound to male and female erythrocytes among thalassemia group was  $95.2 \pm 0.3$  ng/10<sup>6</sup> RBC and  $98.7 \pm 1.3$  ng/10<sup>6</sup> RBC respectively. There was no difference observed in dye binding between male and female erythrocytes from the subjects of both the groups, but AB dye binding to thalassemic erythrocytes (95.1) was significantly less ( $p < 0.05$ ) as compared to binding on erythrocytes of the control (127.9) group.

**Graph 1: ng of AB dye bound to 10<sup>6</sup> RBC in control and Thalassemia Group -Ggender vise**



**Graph 2: ng of AB dye bound to 10<sup>6</sup> RBC in control and Thalassemia Groups**



## Discussion

Glycobiology is the focus of today's research, as this post translational reaction is recognized playing role in the progression of the disease. There is a shift in research interest from metabolic diseases to congenital hemolytic disorders. Where changes in the erythrocyte membranes are not considered to be the primary defect, yet contribute to the pathophysiology of hemolytic process. Alteration in membrane structure of erythrocytes was reported in hemoglobinopathies such as HbS<sup>(21)</sup> G6PDH deficiency, hereditary spherocytosis<sup>(22)</sup> and paroxysmal nocturnal hemoglobinuria (PHN) a Genetic acquired hemolytic disease which show impaired sialylation of glycosphingolipids and metabolic disorder of membrane Glycoconjugates.<sup>(23)</sup>

Sialic acids a component of Glycoconjugates, present on the surface of erythrocyte membrane contribute almost all negative charge,<sup>(24)</sup> thus binding of AB was indirectly used to assess the negative charge on these red cells. Our result indicates there is a significant difference in charge on erythrocytes between the groups. Thalassaemic erythrocyte shows low AB dye binding, indicating less Sialic acid and in turn less negative charge on the membrane.

Our results are supported by existing results on erythrocyte membrane from thalassaemics where, sialic acid (total, protein bound or lipid bound) has shown decrease level. Experimental data also suggest an increase of sialic acid in serum of thalassaemic individuals. Parameters which alter rheological properties like decreased osmotic fragility, increased aggregation, microcytosis, low MCV of red cell also indirectly supports our findings of decreased charge on thalassaemic red cell membrane.

Short life span of thalassaemic erythrocyte could be because of loss of membrane Sialic acid, which mark them old and making them recognizable by RE cell receptor for early clearance<sup>(25)</sup> as desialylation, a negative consequence may appear to protect individual from non-functional red cells.

Erythrocytes in circulation constantly encounter elevated oxygen pressure. Ferrous ion acts as a strong catalyst to cause auto-oxidation stress. Even though defense enzymes of cell counteract this stress, iron overload and deficient antioxidant defense mechanism in thalassaemia cause damage on membrane protein and lipid<sup>(26)</sup> such denatured rafts are shed off from the membrane as a part of cell protection process. Reduced Sialic acid to protein ratio has been reported in beta thalassaemia.<sup>(27)</sup> Oxidative stress was also observed in certain genetic diseases like beta hemoglobinopathies (HbS, thalassaemia) G6PDH deficiency, hereditary spherocytosis, PHN. Due to loss of Sialic acid erythrocytes become more sensitive to osmotic shock and thus leads to enhanced eryptosis, (suicidal death of erythrocytes).<sup>(22)</sup> Accumulation of alpha globin chain at cytoskeleton in beta thalassaemia with loss of thiol group and denaturation of protein 3 & 4.1band designate oxidative damage on membrane.<sup>(28)</sup>

Kahane I et al<sup>(29)</sup> used cationised ferritin and Ruthenium red dye to visualize membrane based Sialyl residue, thalassaemia erythrocyte membrane has exhibited less dense material as compared to normal. We are yet to know the exact molecular mechanism of observed reduction in Sialic acid. It could be defective biogenesis of membrane under genetic defect or early removal of damaged components from membrane.

Whatever may be the molecular mechanism involved in loss of Sialic acid carrying negative charge from thalassaemic erythrocyte, we believe, in addition to a defective beta chain, exposure to oxidative stress could lead to loss of negatively charged Sialic acid augment removal of tinted erythrocyte from circulation

thereby protecting the individual from non-functional red cells but in effect increasing the severity of disease.

Further research towards role of sialyltransferase and sialidase is needed. Also protecting Sialic acid on the erythrocyte membrane or masking of penultimate galactose epitope may provide defensive shield.

## Conclusion

We believe that our data support the perception that there is decreased negative charge in the form of Sialic acid from the erythrocyte membrane in thalassaemia which is reflected by cationic Alcian Blue dye binding.

## Acknowledgment

The authors are thankful to the Indian Council of Medical Research (ICMR), New Delhi for providing financial support in the form of STS 2016 project. The authors deny any conflicts of interest related to this study.

## References

1. Varki, A. Sialic acids in human health and disease, *Trends in Molecular Medicine*; 2008;14(8),351-60.
2. Higgs DR, Thein SL, Wood WG. Distribution and population genetics of the thalassaemias. In: Weatherall DJ, Clegg JB, eds. *The Thalassaemia Syndromes*. 4th ed. Blackwell Science, Oxford 2001;p.237-284
3. Greer JP, Daniel A, Glader B et al, *Thalassaemia and related disorders*. In Greer JP eds. *Wintrobe's Clinical Hematology*, 13<sup>th</sup> ed. Lippincott Williams and Wilkins, Philadelphia. 2014, P1319.
4. Vetter B, Neu-Yilik G, Kohne E, Arnold R, Sinha P, Gaedicke G, Ivancevic V, Kulozik AE. Dominant beta-thalassaemia: a highly unstable hemoglobin is caused by a novel 6 bp deletion of the beta-globin gene. *Br J Haematol*. 2000;108(1):176-81.
5. Sagare Aparna, Dhiraj J Trivedi. Assessment of Transferrin Saturation as an Indicator of Iron Overload in Homozygous & Heterozygous Form of Thalassaemia. *RJPBCS*, 2014,5(1),668-673.
6. Dhiraj J Trivedi, Aparna Sagare. Assessment of Iron Overload in Homozygous and Heterozygous Beta Thalassaemic Children below 5 Years of Age, *JKIMSU*, 2014 3(2),9-14.
7. Onde L de S, Estevo I da F, Rocha MIP. Oxidative stress and antioxidant status in beta-thalassaemia heterozygotes. *Revista Brasileira de Hematologia e Hemoterapia*. 2013;35(6):409-413.
8. Meral A, Tuncel P, Surmen-Gur E, Ozbek R, Ozturk E, Gunay U. Lipid peroxidation and antioxidant status in beta-thalassaemia. *Pediatr Hematol Oncol*. 2000;17(8):687-93.
9. Wilairat P, Kittikalayawong A, Chaicharoen S. The thalassaemic red cell membrane. *Southeast Asian J Trop Med Public Health*, 1992,23 (supl-2), 74-78.
10. R. Advani, S. Sorenson, E. Shinar, W. Lande, E. Rachmilewitz, S.L. Schrier. Characterization and Comparison of the Red Blood Cell Membrane Damage in Severe Human alpha and beta Thalassaemia; 1992; *Blood*, 79(4); 1058-1063.
11. Paul Whiteman. The quantitative measurement of Alcian Blue-glycosaminoglycan complexes; *Biochem J*, 1973;131(2);343-350.

12. Podracka L, Sasinka M, Boor A, Kovacs L. Change in physico chemical characteristics of red blood cells in children with nephrotic syndrome. *International Urology and Nephrology*, 1996,28(4),593-99.
13. Klinger M, Kramarz S, Wendycz D, kopec W. Different erythrocyte and platelet surface electric charge in various types of glomerulonephritis. *Nephrol Dial Transplant*, 1997,12,707-712.
14. Winkel JGJ, Wetzels JFM, Duijnhoven JLP, Koene RPA, Capel PJA. Red blood cell surface charge and alcian blue binding. *Nephrol Dial Transplant*, 1987,2,280-81.
15. Gambaro G, Baggio B, Cicerello E, Mastrosimone S, Marzaro G, Borsatti A, Crepaldi G. Abnormal erythrocyte charge in diabetes mellitus. Link with microalbuminuria. *Diabetes*, 1988,37:6;745-748.
16. Baba Y, Kai M. The lower level of surface electric charge in diabetes mellitus. *Clinica chimica Acta*, 1978,84,247-249.
17. Levin M, Smith C, Walters MDS, Gascoin P, Barratt TM. Steroid responsive nephrotic syndrome: a generalized disorder of membrane negative charge. *Lancet*, 1985, ii, 239-42.
18. Yavuz D, Toprak A, Budak Y, Ersozl HO, Deynelil O, Tezcan H, Akalin S, Urine Glycosaminoglycans excretion in newly diagnosed essential hypertensive patients. *Am Assoc. Cline Chem*, 2000.
19. Shine I and Lal S. A strategy to detect beta thalassemia minor. *Lancet*.1977, 1, 692-94.
20. Modified Betke method for estimation of HbF. *Dacie & Lewis, Practical Haematology*, Elsevier(eds), 11<sup>th</sup> edition
- Riggs MG, Ingram VM 1977, *Biochemistry biophysics res commn* ,1977,74,192-198.
21. Lang KS, Roll B, Myssina S, Schittenhel M. Enhanced erythrocyte apoptosis in sickle cell anemia, thalassemia and G6PD deficiency, *Cell physiol. Biochem*, 2002,12(5-6),365-72.
22. Nakakuma H, Kawaguchi T, Horikawa K. Altered expression of gangliosides in erythrocyte of PHN. *J Clin Invest*, 1990,85(5),1456-1461.
23. Eylar EH, Madoff MA, Brody OV, Oncley JL. The contribution of Sialic acid to the surface charge of the erythrocyte. *J. Biol chem.*, 1962,237,1992-2000.
24. Durocher JR, Payne RC, Conrad ME. Role of Sialic acid in erythrocyte survival. *Blood*, 1975,45(1),11-20.
25. Pavri RS, Baxi AJ, Bhatia HM. Oxidative injury to the red cell in thalassemia and the protective role of antioxidants. *Indian J Med Res*, 1988,88;356-59.
26. Kahane I, Ben-Chetrit E, Shifter A, Rachmilewitz EA. The erythrocyte membrane in beta-thalassemia. Lower Sialic acid levels in glycophorin. *Biochim Biophys Acta*. 1980, 596(1),10-17.
27. R Advani, S Sorenson, E. Shinar, W Lande, E. Rachmilewitz, SL Schrier. Characterization and Comparison of the Red Blood Cell Membrane Damage in Severe Human  $\alpha$  and  $\beta$ -Thalassemia [www.bloodjournal.org](http://www.bloodjournal.org), 2016.
28. I. Kahane, A. Polliack, E. A. Rachmilewitz, E. A. Bayer, E. Skutelsky. Distribution of Sialic acid on the red cell membrane in beta thalassemia. *Nature*, 1978, 271(5646),674-75.